Presto Gel is registered as a medical device, Class I, according to Directive 93/42/EEC



Directions for Use

Presto Gel brings rapid, lasting relief to inflamed, swollen & painful Haemorrhoids

Contains natural active ingredients (no drugs)

Methodology

Presto Gel contains a unique and advanced composition of highly active natural ingredients in a soothing, effective formula that protects inflamed tissues from further bacterial infection and/or contact with faecal acids and toxins, which are the main cause of anal discomforts.

This protection does not destroy the bacteria but instead helps create optimum conditions to support your body's natural biological wound healing mechanisms and allows them to accelerate the normal process of cell generation and tissue repair.

Directions for use

For topical and/or internal rectal application.

- Keep the anal area hygienically clean after every bowel movement by washing with Dermatone tea tree soapless soap or an equivalent low allergy body soap, preferably having mild antibacterial properties. Wash the area at least 2-3 times daily.
- 2. Remove the cap from the tube and use the spike to pierce the underlying seal. Screw on the applicator tip.
- 3. After washing rectal area, use a finger-tip to apply Presto Gel to external haemorrhoids or if only treating internal haemorrhoids, apply to the area needed to lubricate the passage of applicator tip. Carefully insert the applicator to its full length. Squeeze tube to expel a small quantity of the Gel. Withdraw and clean the applicator. Refit cap. Gel may be digitally inserted for easier portion control.
- 4. Apply Presto Gel at least 4-6 times daily to maintain the required antibacterial barrier. A little and often gives best results. Keep finger nails short and clean. Users may experience a temporary, pleasant 'cooling' sensation after each application.

Individual response may vary so do not reduce the rate of application of Presto Gel until your body has had time to bring about the desired results. Thereafter a reduced maintenance program may be followed but is not mandatory.

Application of Presto Gel does not need to be liberal. A quantity about the size of a green garden pea (250 mg) should be adequate to provide the required protective barrier film to local bacteria etc.

Presto and it's over!

Each tube contains 25g which should be sufficient for 100 applications.

Where fissures or tissue damage exists, a brief, burning sensation may occur upon first application of the gel. This sensation typically fades with continued application. Washing the rectal area will stop the sensation but also compromise the protective barrier. This procedure is rarely if ever necessary.

Precautions

Avoid contact with eyes. Do not swallow. Use as directed. If symptoms persist, see your health professional.

Duration of Use

Unlimited.

Suitability

For use by men and women of all ages, all pregnant, postnatal and lactating women and children above the age of 12. The use of this gel for children under the age of 12 must be under a physician's supervision.

Composition

Water, Aloe Barbadensis Extract, Triethanolamine, Carbomer, Geranium Thunbergii Extract, Pogostemon Cablinleaf Extract.

Side Effects

None expected. If an allergic reaction occurs during use, (local irritation and/or rash) please refer to your health professional before continuing use of the gel.

Drug Interactions

None.

Storage

Store in a cool dry place away from children.

For further information, please visit www.prestogel.co.nz

